arXiv:math/0601255v1 [math.GT] 11 Jan 2006

EXTENDING HOMEOMORPHISMS FROM 2-PUNCTURED SURFACES TO HANDLEBODIES

ALESSIA CATTABRIGA AND MICHELE MULAZZANI

Abstract. Let H_g be a genus g handlebody, and $\mathrm{T}_g = \partial\mathrm{H}_g$ a closed connected orientable surface. In this paper we find a finite set of generators for $\mathcal{E}_2^g$, the subgroup of $\mathrm{PMCG}_2(\mathrm{T}_g)$ consisting of the isotopy classes of homeomorphisms of T_g which admit an extension to the handlebody keeping a properly embedded trivial arc fixed. This subgroup turns out to be important for the study of knots in closed 3-manifolds via $(g,1)$-decomposition. In fact, the knots represented by the isotopy classes belonging to the same left cosets of $\mathcal{E}_2^g$ in $\mathrm{PMCG}_2(\mathrm{T}_g)$ are equivalent.

Mathematics Subject Classification 2000: Primary 20F38; Secondary 57M25, 57N10.
Keywords: $(g,1)$-decompositions of knots, mapping class groups, extending homeomorphisms, handlebodies.

1. Introduction

The study of knots and links in closed 3-manifolds via Heegaard decompositions was started by Doll in [6], where the notion of (g,b)-decompositions of links is introduced. In particular, the case of $(1,1)$-decompositions of knots in lens spaces and in the 3-sphere has been extensively investigated in several papers (see for example [2, 3, 4, 7, 8, 9, 10]). More generally, any knot K in a 3-manifold M of Heegaard genus h admits a $(g,1)$-decomposition for a suitable $g \geq h$. This means that there exists a genus g Heegaard splitting $(\mathrm{H}_g, \mathrm{H}'_g)$ of M such that $\mathrm{T}_g = \partial\mathrm{H}_g = \partial\mathrm{H}'_g$ is transverse to K and splits it in two arcs, $A \subset \mathrm{H}_g$ and $A' \subset \mathrm{H}'_g$, parallel to T_g (for details see [5]). As a consequence, the knot K can be represented, up to equivalence, by an element of $\mathrm{PMCG}_2(\mathrm{T}_g)$, the pure mapping class group of the twice punctured surface of genus g. Moreover, if a homeomorphism $\psi : (\partial\mathrm{H}_g, \partial A) \to (\partial\mathrm{H}_g, \partial A)$ admits an extension to the handlebody H_g fixing A, the knot K_ψ, represented by the isotopy class of ψ, is the trivial knot in the connected sum of g-copies of $\mathbf{S}^1 \times \mathbf{S}^2$. Therefore, if we denote with $\mathcal{E}_2^g$ the subgroup of $\mathrm{PMCG}_2(\mathrm{T}_g)$, consisting of the elements which admit an extension to H_g fixing A, and $\mathcal{K}_{g,1}$ is the set of all the knots admitting a $(g,1)$-decomposition, there exists a surjective map

$$\Theta_g : \mathrm{PMCG}_2(\mathrm{T}_g)/\mathcal{E}_2^g \to \mathcal{K}_{g,1},$$

where $\mathrm{PMCG}_2(\mathrm{T}_g)/\mathcal{E}_2^g$ denotes the set of left cosets of $\mathcal{E}_2^g$ in $\mathrm{PMCG}_2(\mathrm{T}_g)$.

The aim of this paper is to find a finite set of generators for $\mathcal{E}_2^g$. In Section 2, we introduce notations, describe some significant elements of $\mathcal{E}_2^g$, which have an explicit topological meaning, and recall some known facts. The main result of the paper is given in Section 3, where a finite set of generators for $\mathcal{E}_2^g$ is obtained, for each $g \geq 1$.

2. Preliminaries and notations

Given a closed orientable connected surface T_g of genus $g \geq 0$ and n distinguished points $P_1, \ldots, P_n$ on it, called *punctures*, we denote with $\mathcal{H}_n(\mathrm{T}_g)$ (resp. $\mathcal{F}_n(\mathrm{T}_g)$) the group of orientation-preserving homeomorphisms $h : \mathrm{T}_g \to \mathrm{T}_g$ such that $h(\{P_1, \ldots, P_n\}) = \{P_1, \ldots, P_n\}$ (resp. $h(P_i) = P_i$ for $i = 1, \ldots, n$). The *n- punctured mapping class group* $\mathrm{MCG}_n(\mathrm{T}_g)$ (resp. the *n-punctured pure mapping class group* $\mathrm{PMCG}_n(\mathrm{T}_g)$) of T_g is the group of the isotopy classes of elements of $\mathcal{H}_n(\mathrm{T}_g)$ (resp. $\mathcal{F}_n(\mathrm{T}_g)$). Clearly, neither $\mathrm{MCG}_n(\mathrm{T}_g)$ nor $\mathrm{PMCG}_n(\mathrm{T}_g)$ depend on the choice of the punctures on T_g. There is an exact sequence

$$1 \to \mathrm{PMCG}_n(\mathrm{T}_g) \to \mathrm{MCG}_n(\mathrm{T}_g) \to \Sigma_n \to 1,$$

where Σ_n is the symmetric group of degree n. Moreover, we denote with $\mathrm{MCG}(\mathrm{T}_g)$ the mapping class group of T_g with no punctures.

Let H_g be an orientable handlebody of genus g and denote with $A \subset \mathrm{H}_g$ a properly embedded trivial arc (i.e., there exists a disk $D \subset \mathrm{H}_g$ with $A \cap D = A \cap \partial D = A$ and $\partial D - A \subset \partial \mathrm{H}_g$). The boundary of H_g is a closed orientable connected surface of genus g with two distinguished punctures P_1 and P_2 on it, which are the endpoints of A. In this article we are interested in determining generators for the subgroup $\mathcal{E}_2^g$ of $\mathrm{PMCG}_2(\mathrm{T}_g)$, consisting of the elements which admit an extension to H_g fixing A.

The model for a genus g handlebody H_g will be the one depicted in Figure 1, and $A \subset \mathrm{H}_g$ will be the properly embedded trivial arc dashed in the figure.

In order to simplify notation, we will use the same symbol to denote a homeomorphism and its isotopy class. Moreover, $\overline{\psi}$ will denote an extension to H_g of a homeomorphism ψ of T_g. The right-handed Dehn twist along a simple closed curve e will be denoted with t_e, while $s_{P,e}$ will denote the spin of P about e (see [1]).

Now we briefly recall some homeomorphisms of T_g that admit an extension to H_g keeping A fixed (for details see [11]).



Figure 1. A model for H_g.

Rotation: let $\overline{\rho}$ be the clockwise $\frac{2\pi}{g}$-rotation of H_g around the axis containing the punctures P_1 and P_2. The restriction of this homeomorphism to T_g will be denoted with ρ.

Semitwist of a knob: let D_i be the 2-cell depicted in Figure 1 and let $\delta_i = \partial D_i$, for $i = 1, \dots, g$. Then D_i cuts away from H_g a solid torus K_i, containing the i-th handle V_i, that will be called the i-th knob. We denote with $\overline{\omega}_i$ the homeomorphism of H_g obtained by cutting H_g along D_i, giving a semitwist counterclockwise to K_i and gluing it back, for $i = 1, \dots, g$. Moreover, we set $\omega_i = (\overline{\omega}_i)_{|\mathrm{T}_g}$. Notice that ω_i^2 is isotopic to the Dehn twist along δ_i.

Twisting a handle: Referring to Figure 1, let $\beta_i = \partial B_i$, for $i = 1, \dots, g$. We denote with τ_i the right-handed Dehn twist along β_i. Note that τ_i admits an extension to H_g whose effect is to give a complete twist to the i-th handle, for $i = 1, \dots, g$.

Slides: Referring to Figure 1, let Z_i (resp. Z_i') be the center of the properly embedded meridian disc $B_i \subset \mathrm{H}_g$ (resp. $B_i' \subset \mathrm{H}_g$). Moreover, denote with H_g^i the genus $(g-1)$ handlebody obtained by removing the i-th handle V_i from H_g, and let e

(resp. e') be a simple closed curve on $\partial \mathrm{H}_g^i - \{P_1, P_2\}$ containing Z_i (resp. Z_i'), with $Z_i' \notin e$ (resp. $Z_i \notin e'$). Such a curve will be called an *i-loop* (resp. an *i'-loop*). Consider the spin $s_{Z_i,e}$ on $\partial \mathrm{H}_g^i$. We can suppose, up to isotopy, that $s_{Z_i,e}(B_i) = B_i$ and $s_{Z_i,e}(B_i') = B_i'$. So, extending $s_{Z_i,e}$ by the identity on ∂V_i, we obtain a homeomorphism of T_g which will be denoted with $\sigma_{i,e}$ and called the *slide* of Z_i along e. In a completely analogous way, given a i'-loop e', we can define a slide of Z_i' along e' and denote it with $\sigma'_{i',e'}$. Let $\mathcal{S} = \{\sigma_{i,e} \mid e \text{ is an } i\text{-loop }, i = 1, \ldots, g\} \cup \{\sigma'_{i',e'} \mid e' \text{ is an } i'\text{-loop }, i' = 1, \ldots, g\}$. Moreover, we set $\theta_{ij} = \sigma_{i,e_{ij}}$, $\xi_{ij} = \sigma_{i,g_{ij}}$, $\eta_{ik} = \sigma_{i,f_{ik}}$ (resp. $\theta'_{ij} = \sigma'_{i',e'_{ij}}$, $\xi'_{ij} = \sigma'_{i',g'_{ij}}$, $\eta'_{ik} = \sigma'_{i',f'_{ik}}$), where $e_{ij}, g_{ij}, f_{ik}, e'_{ij}, g'_{ij}, f'_{ik}$ are the curves depicted in Figure 2, with $i, j = 1, \ldots, g$, $i \neq j$, and $k = 1, 2$.



FIGURE 2. The loops $e_{ij}, g_{ij}, f_{ik}, e'_{ij}, g'_{ij}, f'_{ik}$, on H_g^i.

Exchanging two knobs: referring to Figure 3, let C_i (resp. C_j) be the 2-cell on ∂H_g^i (resp. ∂H_g^j) bounded by ν_i (resp. ν_j) not containing K_j (resp. K_i). Denote with D a tubolar neighborhood of $C_i \cup e \cup C_j$ on T_g. Let ρ'_{ij} be the homeomorphism of T_g obtained by exchanging ∂K_i with ∂K_j in such a way that ρ'_{ij} twists $C_i \cup e \cup C_j$ inside D through π radians in the clockwise direction. So, $\rho'_{ij}(C_i) = C_j$, $\rho'_{ij}(C_j) = C_i$ $\rho'_{ij}(e) = e$ and ρ'_{ij} is the identity outside a tubolar neighborhood of $K_i \cup e \cup K_j$. We set $\rho_{ij} = \omega_j \omega_i \rho'_{ij}$, for $i, j = 1, \ldots, g$ and $i \neq j$.



Figure 3. Exchanging two knobs.

Proposition 1 ([11]). *Let* $e_0, \ldots, e_n$ *(resp.* $e'_0, \ldots, e'_n$*) be i-loops (resp. i′-loops) such that* e_0 *(resp.* e'_0*) is homotopic to* $e_1 \cdots e_n$ *(resp.* $e'_1 \cdots e'_n$*) on* $\partial \mathrm{H}^i_g - \{P_1, P_2, Z'_i\}$ *rel* Z_i *(resp.* $\partial \mathrm{H}^i_g - \{P_1, P_2, Z_i\}$ *rel* Z'_i*). Then* σ_{i,e_0} *(resp.* σ'_{i',e'_0}*) is isotopic to* $\sigma_{i,e_1} \cdots \sigma_{i,e_n}$ *(resp.* $\sigma'_{i',e'_1} \cdots \sigma'_{i',e'_n}$*) modulo* τ_i*.*

Proposition 2 ([11]). *The subgroup* $\mathcal{E}^g$ *of* $\mathrm{MCG}(\mathrm{T}_g)$*, containing the elements of* $\mathrm{MCG}(\mathrm{T}_g)$ *that extend to* H_g *is generated by* $\rho, \rho_{12}, \xi_{12}, \theta_{12}, \tau_1, \omega_1$*.*

3. Main result

In this paragraph we establish the main result of the article.

Theorem 3. *The subgroup* $\mathcal{E}^g_2$ *of* $\mathrm{PMCG}_2(\mathrm{T}_g)$ *is generated by* $\rho, \rho_{12}, \xi_{12}, \theta_{12}, \tau_1, \omega_1, \eta_{11}, \eta_{12}$ *if* $g > 1$*, and* $\tau_1, \omega_1, t_\gamma$ *if* $g = 1$*, where* γ *is the curve depicted in Figure 4.*

Before giving the proof we need some preparatory lemmas. Denote with $\mathcal{Y}^g$ the subgroup of $\mathcal{E}^g_2$ generated by $\rho_{ij}, \xi'_{ij}, \xi_{ij}, \theta'_{ij}, \theta_{ij}, \tau_i, \omega_i, \eta_{i1}, \eta_{i2}, \eta'_{i1}, \eta'_{i2}$, with $i, j = 1, \ldots, g$ and $i \neq j$.

Lemma 4. *Let* e *(resp.* e'*) be an i-loop (resp. i′-loop). Then* $\sigma_{i,e}$ *(resp.* $\sigma'_{i,e'}$*) is isotopic to a product of* $\xi_{ij}, \theta_{ij}, \eta_{ik}, \tau_i$ *(resp.* $\xi'_{ij}, \theta'_{ij}, \eta'_{ik}, \tau_i$*), for* $j = 1, \ldots, g$*,* $k = 1, 2$ *and* $i \neq j$*.*

Proof. Since e_{ij}, g_{ij}, f_{ik}, (resp. $e'_{ij}, g'_{ij}, f'_{ik}$) with $j = 1, \ldots, n$, $k = 1, 2$ and $i \neq j$, are a free set of generators of $\pi_1(\partial \mathrm{H}^i_g - \{P_1, P_2, Z'_i\}, Z_i)$, (resp. $\pi_1(\partial \mathrm{H}^i_g - \{P_1, P_2, Z_i\}, Z'_i)$) the lemma follows from Proposition 1. □

Lemma 5. *For each $f \in \mathcal{E}_2^g$ there exists an element $f_0 \in \mathcal{Y}_g$ such that $\overline{f}_0\overline{f}$ is the identity on the i-th handle.*

Proof. By Lemma 4, $\mathcal{S} \subseteq \mathcal{Y}_g$. Therefore, using the same arguments as in the proof of [11, Lemma 4.4] we have that $\overline{f}_0\overline{f}$ is the identity on a meridian disk of the i-th handle and so, up to isotopy, on all the handle. $\square$

Proposition 6. *For $g \geq 1$ and $1 \leq m < n$, consider the homomorphism $j_{g,n,m} : \mathrm{PMCG}_n(\mathrm{T}_g) \to \mathrm{PMCG}_m(\mathrm{T}_g)$ induced by the inclusion. Then $\ker j_{g,n,m} \cong \pi_1(F_{m,n-m}(\mathrm{T}_g))$, where $F_{m,n-m}(\mathrm{T}_g)$ denotes the configuration space of $n-m$ points in $\mathrm{T}_g - \{P_1, \dots P_m\}$.*

Proof. Let $P_1, \dots, P_n$ be distinct points on T_g. As in [1, Theorem 4.1], it is possible to prove that the evaluation map $\epsilon : \mathcal{F}_m(\mathrm{T}_g) \to F_{m,n-m}(\mathrm{T}_g)$, given by $\epsilon(\psi) = (\psi(P_{m+1}), \dots, \psi(P_n))$, is a locally trivial fibering with fibre $\mathcal{F}_n(\mathrm{T}_g)$, so the following exact sequence holds:

$$\pi_1(F_{m,n-m}(\mathrm{T}_g), (P_{m+1}, \dots, P_n)) \xrightarrow{d_\#} \pi_0(\mathcal{F}_n(\mathrm{T}_g), id) \xrightarrow{j_\#} \pi_0(\mathcal{F}_m(\mathrm{T}_g), id) \to 1.$$

Since $\pi_0(\mathcal{F}_k(\mathrm{T}_g), id) = \mathrm{PMCG}_k(\mathrm{T}_g)$ and $j_\#$ is induced by the inclusion, it is enough to prove that $\ker(d_\#) = 1$ to get the statement. Proceeding as in the proof of [1, Lemma 4.2.1], we obtain that $\ker(d_\#)$ is contained in the center of $\pi_1(F_{m,n-m}(\mathrm{T}_g), (P_{m+1}, \dots, P_n))$. Now we will prove, by induction on k, that the center of $\pi_1(F_{m,k}(\mathrm{T}_g), *)$ is trivial. For $k = 1$ we have $\pi_1(F_{m,1}(\mathrm{T}_g), *) \cong \pi_1(\mathrm{T}_g - \{P_1, \dots, P_m\}, *)$, which is a free group on $2g + m - 1 > 1$ generators. Now let $k > 1$. By [1, Theorem 1.2] for each $1 \leq r < k$ the projection $\pi : F_{m,k}(\mathrm{T}_g) \to F_{m,r}(\mathrm{T}_g)$ is a locally trivial fibering with fiber $F_{m+r,k-r}(\mathrm{T}_g)$. Since $\pi_3(F_{m,1}(\mathrm{T}_g)) = \pi_3(\mathrm{T}_g - \{P_1, \dots, P_m\}) = 1$ and $\pi_2(F_{m,1}(\mathrm{T}_g)) = \pi_2(\mathrm{T}_g - \{P_1, \dots, P_m\}) = 1$, from the long exact sequence of the above fibration, for $r = 1$, we get $\pi_2(F_{m+1,k-1}(\mathrm{T}_g)) \cong \pi_2(F_{m,k}(\mathrm{T}_g))$. So an inductive argument shows that $\pi_2(F_{m,k-1}(\mathrm{T}_g)) \cong \pi_2(F_{m+1,k-2}(\mathrm{T}_g)) \cong \cdots \cong \pi_2(F_{m+k-2,1}(\mathrm{T}_g)) = 1$. Then, from the long exact sequence of the fibration in the case $r = k-1$ we get

$$1 \to \pi_1(F_{m+k-1,1}(\mathrm{T}_g)) \xrightarrow{i_\#} \pi_1(F_{m,k}(\mathrm{T}_g)) \xrightarrow{\pi_\#} \pi_1(F_{m,k-1}(\mathrm{T}_g)) \to 1.$$

Now, by the induction hypothesis, the center of $\pi_1(F_{m,k-1}(\mathrm{T}_g))$ is trivial. So the center of $\pi_1(F_{m,k}(\mathrm{T}_g))$ is contained in $\ker(\pi_\#) \cong i_\#(\pi_1(F_{m+k-1,1}(\mathrm{T}_g))) \cong \pi_1(F_{m+k-1,1}(\mathrm{T}_g))$, which is centerless. $\square$

Remark 7. As in [1, pp. 158-160], a system of generators of $\ker j_{g,n,m}$ can be constructed as follows. Let E_n be a set of generators of $\pi_1(\mathrm{T}_g - \{P_1, \dots P_m\}, P_n)$, such that for each $e \in E_n$, we have

$e \cap \{P_1, \dots, P_n\} = \{P_n\}$. Moreover, for each $i = m+1, \dots, n-1$, denote with E_i a set of generators of $\pi_1(\mathrm{T}_g - \{P_1, \dots P_m\}, P_i)$ such that $e \cap \{P_1, \dots, P_n\} = \{P_i\}$ and each loop of E_i is free homotopic on $\mathrm{T}_g - \{P_1, \dots, P_m\}$ to a loop in E_n. Then $\ker j_{g,n,m}$ is generated by $\{s_{i,e} \,|\, e \in E_i, i = m+1, \dots, n\}$.



Figure 4. The case of genus one.

Now we are ready to prove Theorem 3.

Proof. Suppose that $\mathcal{Y}^g = \mathcal{E}_2^g$. From the relations in [11, Section 3], and since $\eta_{ik} = \rho^{i-1}\eta_{1k}\rho^{-(i-1)}$, $\eta'_{i1} = \omega_i^{-1}\eta_{i1}\omega_i$ and $\eta'_{i2} = \omega_i\eta_{i2}\omega_i^{-1}$, for $i = 1, \dots, g$, the subgroup $\mathcal{Y}^g$ is generated by $\rho, \rho_{12}, \xi_{12}, \theta_{12}, \tau_1, \omega_1, \eta_{11}, \eta_{12}$ if $g > 1$, and by $\tau_1, \omega_1, \eta_{11}, \eta_{12}$ if $g = 1$. This completes the proof when $g > 1$. If $g = 1$, referring to Figure 4, we have $\eta_{11} = t_{\varphi_1}t_{\varphi_2}^{-1} = t_{\varphi_1}$ and $\eta_{12} = t_{\chi_1}t_{\chi_2}^{-1} = t_{\chi_1}$. Moreover, $\varphi_1 \simeq \gamma$, $\chi_1 \simeq \varphi'$ and $\omega_1(\varphi_1) = \varphi'$.

Now we show, by induction on the genus, that $\mathcal{Y}^g = \mathcal{E}_2^g$. Let $\widehat{\mathcal{E}}_2^g$ be the subgroup of $\mathcal{E}_2^g$ containing the isotopy classes of the homeomorphisms that are the identity on the boundary of the g-th handle V_g. By Lemma 5, up to composing with an element of $\mathcal{Y}^g$, we can suppose that each $f \in \mathcal{E}_2^g$ belongs to $\widehat{\mathcal{E}}_2^g$. So it is enough to show that $\widehat{\mathcal{E}}_2^g \subseteq \mathcal{Y}^g$.

Let $f : \mathrm{T}_g \to \mathrm{T}_g$ be a homeomorphism fixing ∂V_g pointwise and whose isotopy class belongs to $\widehat{\mathcal{E}}_2^g$. By cutting out the g-th handle, and capping with the two disks B_g and B'_g, we can identify f with a homeomorphism f' of T_{g-1}, such that $f'_{|_{B_g \cup B'_g}} = \mathrm{Id}$ and $f' = f$ on $\mathrm{T}_g - (B_g \cup B'_g)$. Moreover, by shrinking B_g and B'_g to their centers

Z_g and Z'_g, the map f' becomes a map $\widetilde{f}$ of T_{g-1} fixing Z_g and Z'_g. In order to simplify the notation, we set $P_3 = Z_g$ and $P_4 = Z'_g$. Obviously, $\widetilde{f}$ extends to H_{g-1} fixing A. Note that isotopies of $\widetilde{f}$ rel $\{P_1, P_2, P_3, P_4\}$ correspond to isotopies of f rel $\{P_1, P_2\}$ and the isotopy class of $\widetilde{f}$ is trivial in $\mathrm{PMCG}_4(\mathrm{T}_{g-1})$ if and only if the isotopy class of f belongs to the subgroup generated by τ_g in $\mathrm{PMCG}_2(\mathrm{T}_g)$.

Let $\widetilde{\mathcal{E}}^{g-1}$ be the subgroup of $\mathrm{PMCG}_4(\mathrm{T}_{g-1})$ consisting of the elements which extend to the handlebody keeping A fixed. Note that for each $g \in \widetilde{\mathcal{E}}^{g-1}$ we have $g = \widetilde{f}$, for a suitable f. Let $g = 1$. We have that



Figure 5. Spin s_{ij} on $\mathbf{S}^2$.

$\mathrm{PMCG}_4(\mathbf{S}^2) = F_{4,0}/\mathrm{center}(F_{4,0})$ and a set of generators for $\mathrm{PMCG}_4(\mathbf{S}^2)$ is given by $\{s_{ij} \mid i, j = 1, \ldots, 4; i < j\}$, where s_{ij} is the spin of P_j along the curve d_{ij} depicted in Figure 5 (see [1, Lemma 1.8.2, Theorem 4.2]), which is equal to the Dehn twist along the curve d'_{ij}. Obviously, $t_{d'_{ij}}$ extends to the 3-ball fixing A, and therefore $\mathrm{PMCG}_4(\mathbf{S}^2) = \widetilde{\mathcal{E}}^0$. It is easy to see that $s_{14} = \widetilde{\eta'_{11}}$, $s_{13} = \widetilde{\eta_{11}}$, $s_{24} = \widetilde{\eta'_{12}}$, $s_{23} = \widetilde{\eta_{12}}$ and, since d'_{12} is isotopic to d'_{34} on $\mathbf{S}^2$ rel $\{P_1, P_2, P_3, P_4\}$, we have $s_{12} = s_{34} = \widetilde{t_{\delta_1}} = \widetilde{\omega_1^2}$. So the isotopy class of f is generated by $\eta'_{11}, \eta_{11}, \eta'_{12}, \eta_{12}, \omega_1^2, \tau_1$. As a consequence $\widehat{\mathcal{E}}_2^1 \subseteq \mathcal{Y}^1$.

Now let $g > 1$. The homomorphism $j_{g-1,4,2} : \mathrm{PMCG}_4(\mathrm{T}_{g-1}) \to \mathrm{PMCG}_2(\mathrm{T}_{g-1})$ of Proposition 6 restricts to a homomorphism $j = (j_{g-1,4,2})_{|\widetilde{\mathcal{E}}^{g-1}} : \widetilde{\mathcal{E}}^{g-1} \to \mathcal{E}_2^{g-1}$. So a set of generators of $\widetilde{\mathcal{E}}^{g-1}$ is given by the generators of $\ker j$ and the preimages of the generators of $\mathcal{E}_2^{g-1}$. By Remark 7, $\ker j$ is generated by spins of Z_g and Z'_g about appropriate loops not containing P_1 and P_2. So they correspond to slides of Z_g and Z'_g on T_g, which are elements of $\mathcal{Y}_g$ by Lemma 4. Moreover, by the induction hypothesis $\mathcal{E}_2^{g-1} = \mathcal{Y}^{g-1}$. Since we can suppose that the generators of $\mathcal{Y}^{g-1}$ keep Z_g and Z'_g fixed, they correspond to element of $\mathcal{Y}^g$. So, as above, the isotopy class of f is generated by

$\rho_{ij}, \xi'_{ij}, \xi_{ij}, \theta'_{ij}, \theta_{ij}, \tau_i, \omega_i, \eta_{i1}, \eta_{i2}, \eta'_{i1}, \eta'_{i2}$, with $i, j = 1, \ldots, g-1$, $i \neq j$, and $\xi'_{gj}, \xi_{gj}, \theta'_{gj}, \theta_{gj}, \eta_{g1}, \eta_{g2}, \eta'_{g1}, \eta'_{g2}, \tau_g$, with $j = 1, \ldots, g-1$. As a consequence $\widehat{\mathcal{E}}_2^g \subseteq \mathcal{Y}^g$. □

Acknowledgements

Work performed under the auspices of the G.N.S.A.G.A. of I.N.d.A.M. (Italy) and the University of Bologna, funds for selected research topics.

References

[1] J. S. Birman, *Braids, Links, and Mapping Class Groups*, Princeton University Press, Princeton-New Jersey, (1974).

[2] A. Cattabriga and M. Mulazzani, *Strongly-cyclic branched coverings of* $(1,1)$*-knots and cyclic presentations of groups*, Math. Proc. Cambridge Philos. Soc. **135** (2003), 137-146.

[3] A. Cattabriga and M. Mulazzani, $(1,1)$*-knots via the mapping class group of the twice punctured torus*, Adv. Geom. **4** (2004), 263-277.

[4] D. H. Choi and K. H. Ko, *Parametrizations of 1-bridge torus knots*, J. Knot Theory Ramifications **12** (2003), 463-491.

[5] P. Cristofori, M. Mulazzani and A. Vesnin, *Strongly-cyclic branched coverings of knots via* $(g,1)$*-decompositions*, preprint, 2005, arXiv: math.GT/0402393.

[6] H. Doll, *A generalized bridge number for links in 3-manifold*, Math. Ann. **294** (1992), 701-717.

[7] H. Goda, C. Hayashi and H.-J. Song, *A criterion for satellite 1-genus 1-bridge knots*, Proc. Amer. Math. Soc. **132** (2004), 3449-3456.

[8] H. Goda, H. Matsuda and T. Morifuji, *Knot Floer homology of* $(1,1)$*-knots*, Geom. Dedicata **112** (2005), 197-214.

[9] C. Hayashi, *1-genus 1-bridge splittings for knots*, Osaka J. Math. **41** (2004), 371-426.

[10] M. Sakuma, *The topology, geometry and algebra of unknotting tunnels. Knot theory and its applications*, Chaos Solitons Fractals **9** (1998), 739–748.

[11] S. Suzuki, *On homeomorphisms of a 3-dimensional handlebody*, Can. J. Math. **29** (1977), 111-124.

ALESSIA CATTABRIGA, Department of Mathematics, University of Bologna, Piazza di Porta S. Donato, 5, 40126, Bologna (Italy). E-mail: cattabri@dm.unibo.it

MICHELE MULAZZANI, Department of Mathematics, C.I.R.A.M., University of Bologna, Piazza di Porta S. Donato, 5, 40126, Bologna (Italy). E-mail: mulazza@dm.unibo.it